

Cancer
Deaths and Death Rates by Race/Ethnicity
San Bernardino County Residents, 1993-2002

		Year									
		1993	1994	1995	1996	1997	1998	1999	2000	2001	2002
Race Group											
White	Deaths	1,790	1,859	1,758	1,742	1,748	1,724	1,740	1,815	1,895	1,809
	Rate	197.2	205.4	194.3	193.3	194.1	190.7	191.8	199.4	207.1	196.8
Black	Deaths	129	157	167	169	153	170	181	180	190	200
	Rate	102.1	121.5	126.2	125.2	110.5	118.9	122.7	118.1	121.0	123.8
Hispanic	Deaths	269	224	295	251	274	308	333	338	372	410
	Rate	61.0	49.6	63.7	53.1	55.7	59.9	61.9	60.0	62.7	65.8
American Indian	Deaths	3	10	5	5	10	8	8	8	6	10
	Rate	28.6	95.6	47.9	48.0	93.9	73.6	72.2	70.9	52.2	85.4
Asian/Pacific Islander	Deaths	47	55	53	52	60	82	55	81	83	74
	Rate	70.5	79.9	74.5	71.1	79.3	102.5	64.9	90.3	87.2	73.4
All Races	Deaths	2,238	2,305	2,279	2,219	2,245	2,292	2,317	2,424	2,547	2,503
	Rate	144.2	147.2	144.1	139.3	138.8	138.7	137.2	140.3	143.8	137.8

RATES ARE UNRELIABLE WHERE LESS THAN 20 DEATHS OCCURRED

Notes:
(1) White, Black, American Indian, and Asian/Pacific Islander exclude Hispanic ethnicity. Hispanic includes any race category.
(2) Rates are per 100,000 population. (3) ICD-9 Codes 140-208; ICD-10 Codes C00-C97.
(3) Cause of death coding was done using ICD-10 beginning in 1999. Care must be taken in analyzing trend data, as the introduction of a new revision of the ICD can cause major discontinuities.

Sources:
(1) CA Department of Health Services: Death Statistical Master Files, 1993-2002
(2) CA Department of Finance: Population Estimates with Age, Sex, and Race/Ethnic Detail, December 1998
Compiled by San Bernardino County Public Health Department, PAS.